


KR-1W

Wave-Front Analyzer



Perfection for Professionals: KR-1W

Topcon, with its wealth of experience in designing and manufacturing refractometers and other diagnostic equipment for over 50 years, introduces a new diagnostic tool with 5 functions to support the evaluation of the visual performance of the human eye. The KR-1W combines aberrometry, topography, keratometry, pupillometry and autorefractometry in one unit that is unparalleled in terms of functionality and reproducibility. At the same time the KR-1W is extremely easy to operate and allows fast patient throughput. The KR-1W will become the professional's choice for comprehensive diagnosis of many ocular conditions.




Features

- » 5 functions in 1 machine:
(Auto-Refractometry, Keratometry, Aberrometry, Topography and Pupillometry)
- » Multiple maps for overview analysis
- » Decision support for cataract and refractive procedures
- » Less stress with invisible light measurement at topography
- » Easier operation with R / L fully automated measurement and touch panel




A Picture Says More Than a Thousand Words

The multiple maps provide helpful insight for refractive procedures that are done on the corneal surface or intraocularly. It allows an objective quality control of the surgical procedures.




Multi Maps




IOL Selection Maps




Summary Maps
(Continuous Measurement Maps)




Pupillometry Maps




Ocular Maps




PSF/MTF Maps




Corneal Maps



Component Maps



Zernike Vector Maps



Corneal R/L Maps

Refraction Data									
	S	C	A	S	C	A			
1	2.50	-1.25	82	3.25	-1.00	89			
2	2.50	-1.25	82	3.25	-1.00	87			
3	2.50	-1.25	82	3.25	-1.00	88			
4	2.50	-1.25	81	3.00	-1.00	85			
5	2.50	-1.25	80	3.00	-0.75	86			
6									
7									
8									
9									
10									
Ave	2.50	-1.25	81	3.25	-1.00	87			

Refraction-Keratometry

Perfection for Professionals: KR-1W

Topcon, with its wealth of experience in designing and manufacturing refractometers and other diagnostic equipment for over 50 years, introduces a new diagnostic tool with 5 functions to support the evaluation of the visual performance of the human eye. The KR-1W combines aberrometry, topography, keratometry, pupillometry and autorefractometry in one unit that is unparalleled in terms of functionality and reproducibility. At the same time the KR-1W is extremely easy to operate and allows fast patient throughput. The KR-1W will become the professional's choice for comprehensive diagnosis of many ocular conditions.




Features


- » 5 functions in 1 machine:
(Auto-Refractometry, Keratometry, Aberrometry, Topography and Pupillometry)
- » Multiple maps for overview analysis
- » Decision support for cataract and refractive procedures
- » Less stress with invisible light measurement at topography
- » Easier operation with R / L fully automated measurement and touch panel

A Picture Says More Than a Thousand Words

The multiple maps provide helpful insight for refractive procedures that are done on the corneal surface or intraocularly. It allows an objective quality control of the surgical procedures.




Multi Maps




IOL Selection Maps




Summary Maps
(Continuous Measurement Maps)




Pupillometry Maps




Ocular Maps




PSF/MTF Maps




Corneal Maps




Component Maps



Zernike Vector Maps



Corneal R/L Maps



Refraction-Keratometry





Work Flow

The KR-1W provides you with the ideal tool to diagnose, follow up and make treatment plans for a broad range of ocular conditions. It can be very effectively used in your practice to speed up your workflow, improve your communication with patients and to monitor the outcomes of your treatments. With the KR-1W you are able to assess your patients functional visual performance over time or pre / post operatively with unsurpassed reproducibility and reliability.

Measure

Unsurpassed reproducibility and reliability with fully automated measurement and touch panel manipulation make your diagnosis and treatment decision easy.

Diagnosis / Consultation


Enrich your communication with patients showing them the effect of their ocular condition on their visual performance.

Treatment / Surgery

Ideal tool to make treatment plans for a broad range of ocular conditions.

Follow-Up

This work flow will increase patient satisfaction and confidence in their choice of treatment as you can objectively show the patient the outcome of your treatments.





Work Flow

The KR-1W provides you with the ideal tool to diagnose, follow up and make treatment plans for a broad range of ocular conditions. It can be very effectively used in your practice to speed up your workflow, improve your communication with patients and to monitor the outcomes of your treatments. With the KR-1W you are able to assess your patients functional visual performance over time or pre / post operatively with unsurpassed reproducibility and reliability.

Measure

Unsurpassed reproducibility and reliability with fully automated measurement and touch panel manipulation make your diagnosis and treatment decision easy.

Diagnosis / Consultation


Enrich your communication with patients showing them the effect of their ocular condition on their visual performance.

Treatment / Surgery

Ideal tool to make treatment plans for a broad range of ocular conditions.

Follow-Up



This work flow will increase patient satisfaction and confidence in their choice of treatment as you can objectively show the patient the outcome of your treatments.



Reliable Decision Support for the Demanding Cataract & Refractive Surgeon

» IOL Selection

Important factors in making the right IOL selection for your patient at one glance.



※ This thresholds are recommendation from Naoyuki Maeda, MD Osaka University Hospital

1 Corneal Higher Order Aberration Index

This index provides information on the potential visual outcome considering the aberration that cannot be corrected by an IOL. Additional treatments, i.e. contact lens fitting or corneal surface treatments, might be necessary to optimize the outcome.

2 Post Refractive Surgery Index

This index provides information on corneal irregularities possibly caused by previous surface treatments of the cornea. This might lead to the necessity of calculating the power of the IOL with non-standard formulas with special attention.


3 Corneal Spherical Aberration Index

This index provides very useful information on the asphericity of the cornea. This gives you the opportunity to select the right monofocal IOL to compensate for the spherical aberration of the patients' cornea.

4 Corneal Astigmatism Index

This simple display of corneal astigmatism provides information for two important decisions.

- a. Amount of toric correction needed in a toric IOL
- b. Patients with a high corneal astigmatism might not be the right candidate for a multifocal IOL



※Data: provided by Tokyo Medical and Dental University Hospital Facility of Medicine

» Pupillometry

Screening application to evaluate eyes for multifocal IOL implantation or refractive surgery.



Corneal Refractive Surgery


For any refractive procedure it is vitally important to diagnose the pupil very carefully in different light conditions.

Also for refractive surgery this tool gives you the necessary information to plan your treatment.



Dynamic Pupil Diameter Measurement and Pupil Center Determination

This information is very important for premium IOL implantation, as it will give you the possibility to select the right multifocal IOL design for the individual eye and also help exclude extreme cases of pupil decentration before surgery.



 Ocular HOA Map
at Scotopic

 Ocular HOA Map
at Photopic

The index of Ocular Total Higher Order aberration, refraction data are provided in photopic and scotopic condition.

Continuous Measurement Function

Up to 10 measurements can be done continuously to observe the change in ocular HOA during about 10 seconds. This may have implication for dry eye diagnosis in the future.




※Data: provided by Tohoku University

Reliable Decision Support for the Demanding Cataract & Refractive Surgeon

» IOL Selection

Important factors in making the right IOL selection for your patient at one glance.



1 Corneal Higher Order Aberration Index

This index provides information on the potential visual outcome considering the aberration that cannot be corrected by an IOL. Additional treatments, i.e. contact lens fitting or corneal surface treatments, might be necessary to optimize the outcome.

3 Corneal Spherical Aberration Index

This index provides very useful information on the asphericity of the cornea. This gives you the opportunity to select the right monofocal IOL to compensate for the spherical aberration of the patients' cornea.


2 Post Refractive Surgery Index

This index provides information on corneal irregularities possibly caused by previous surface treatments of the cornea. This might lead to the necessity of calculating the power of the IOL with non-standard formulas with special attention.

4 Corneal Astigmatism Index

This simple display of corneal astigmatism provides information for two important decisions.

- Amount of toric correction needed in a toric IOL
- Patients with a high corneal astigmatism might not be the right candidate for a multifocal IOL



※Data: provided by Tokyo Medical and Dental University Hospital Faculty of Medicine

5 Review after Toric IOL Implantation

The result after implantation can be reviewed to check residual ocular astigmatism.

[Go to page 8](#)

» Pupillometry

Screening application to evaluate eyes for multifocal IOL implantation or refractive surgery.




Corneal Refractive Surgery

For any refractive procedure it is vitally important to diagnose the pupil very carefully in different light conditions. Also for refractive surgery this tool gives you the necessary information to plan your treatment.

Dynamic Pupil Diameter Measurement and Pupil Center Determination

This information is very important for premium IOL implantation, as it will give you the possibility to select the right multifocal IOL design for the individual eye and also help exclude extreme cases of pupil decentration before surgery.



Ocular HOA Map at Scotopic

Ocular HOA Map at Photopic

The index of Ocular Total Higher Order aberration, refraction data are provided in photopic and scotopic condition.

Continuous Measurement Function

Up to 10 measurements can be done continuously to observe the change in ocular HOA during about 10 seconds. This may have implication for dry eye diagnosis in the future.




※Data: provided by Tohoku University

Full Flexibility for Your Data Viewing or Data Storage Needs

System Chart


The KR-1W is adaptable to your networking and data filing needs since various combinations are available. The patient database is provided with the optional KR-1W Viewer™ software.



* Contact Topcon subsidiaries or dealers for system configurations.

Precise Data Leads to Precise Diagnosis


Case Report



Clinical Data ; Osaka University Hospital
Editorial supervisor ; Naoyuki Maeda, MD

Keratoconus


This is a case of moderate keratoconus. The results of corneal topography are shown at the top row. Axial Power map ➊ indicates typical topographic appearance of keratoconus such as inferior steepening (warmer colors) and skewed axis. The map for the corneal HOA (higher-order aberration) ➋ shows remarkable vertical coma pattern with advanced wavefront (warmer colors) at the superior portion and delayed wavefront (cooler colors) at the inferior cornea. The second row represents the results of ocular aberrations. The ocular total aberration map ➌ shows delayed wavefront in the center. This suggests that uncorrected visual acuity is not good due to myopic astigmatism. The map for ocular HOA ➍ is similar to that for corneal HOA indicating that ocular HOA are mainly attributing from cornea and best spectacle corrected visual acuity is not good. Simulated retinal images of the Landolt rings ➎ suggests that comet-like image will be seen because of vertical coma. In the lower half of the print, IOL selection maps and four outputs are shown. The output for corneal astigmatism ➏ indicates the existence of significant irregular astigmatism with red signal. The difference between central power and K reading ➐ suggests that conventional IOL calculation might induce refractive error due to topographic abnormality. The output for corneal spherical aberration (SA) suggests that the spherical IOL (positive SA) is more appropriate than aspherical IOL (negative SA) because of the negative corneal spherical aberration ➑. The last output is for corneal cylinder and shows that regular astigmatism is extremely high ➒.




Clinical Data ; Osaka University Hospital
Editorial supervisor ; Takashi Fujikado, MD

Cortical Cataract

This is a case of a 62 year old female with cortical cataract. Her visual acuity is 0.8 with correction in the right eye. The corneal HOA (1) is within normal limit but the Total Ocular HOA (2) shows a higher RMS (red indication) at 6mm pupil diameter. Referring to the Component Maps in the lower half of the print, the coma aberration is higher in oculus than in cornea. It is therefore easily observed that the coma aberration is mainly caused by internal components (crystalline lens) (3). The spherical aberration is also higher in oculus than in cornea, indicating that spherical aberration is increased in the crystalline lens (4). Referring to the Landolt's ring simulation (5), it can be seen that the images are vertically distorted, which show the visual effect of the coma aberration. The image simulation is consisted of ocular HOA only, so it should be noted that this simulated image might not coincide with actual patient vision because the opacity of crystalline lens and subsequent light scattering affects the retinal image differently with ocular HOA.



Slit Lamp Anterior Image




Clinical Data ; Tokyo Medical and Dental University Hospital Faculty of Medicine
Editorial supervisor ; Hiroko Bissen-Miyajima, MD

Pseudophakic Eye with Toric IOL

This measurement shows in the upper half the IOL selection map, where it can be shown that the toric IOL is properly implanted and the corneal astigmatism is compensated by the implant. Reference values parameters are: Corneal Astigmatism (showing -1.79D@86), Ocular Astigmatism (showing -0.07D@91) and Internal Astigmatism (showing -1.72D@176). **1** In the lower half the PSF/MTF maps show that objective visual quality should be good as can be seen from the simulated PSF, that gives an indication of optical image quality, and from the MTF that gives an indication of contrast sensitivity **2**.

Customizable Print Layout




Example

2 analysis maps of your choice can be assembled to print layout on 1 page for printing and / or sending images to external systems, like IMAGENet.™

Precise Data Leads to Precise Diagnosis


Case Report



Clinical Data ; Osaka University Hospital
Editorial supervisor ; Naoyuki Maeda, MD

Keratoconus


This is a case of moderate keratoconus. The results of corneal topography are shown at the top row. Axial Power map ➊ indicates typical topographic appearance of keratoconus such as inferior steepening (warmer colors) and skewed axis. The map for the corneal HOA (higher-order aberration) ➋ shows remarkable vertical coma pattern with advanced wavefront (warmer colors) at the superior portion and delayed wavefront (cooler colors) at the inferior cornea. The second row represents the results of ocular aberrations. The ocular total aberration map ➌ shows delayed wavefront in the center. This suggests that uncorrected visual acuity is not good due to myopic astigmatism. The map for ocular HOA ➍ is similar to that for corneal HOA indicating that ocular HOA are mainly attributing from cornea and best spectacle corrected visual acuity is not good. Simulated retinal images of the Landolt rings ➎ suggests that comet-like image will be seen because of vertical coma. In the lower half of the print, IOL selection maps and four outputs are shown. The output for corneal astigmatism ➏ indicates the existence of significant irregular astigmatism with red signal. The difference between central power and K reading ➐ suggests that conventional IOL calculation might induce refractive error due to topographic abnormality. The output for corneal spherical aberration (SA) suggests that the spherical IOL (positive SA) is more appropriate than aspherical IOL (negative SA) because of the negative corneal spherical aberration ➑. The last output is for corneal cylinder and shows that regular astigmatism is extremely high ➒.




Clinical Data ; Osaka University Hospital
Editorial supervisor ; Takashi Fujikado, MD

Cortical Cataract

This is a case of a 62 year old female with cortical cataract. Her visual acuity is 0.8 with correction in the right eye. The corneal HOA (1) is within normal limit but the Total Ocular HOA (2) shows a higher RMS (red indication) at 6mm pupil diameter. Referring to the Component Maps in the lower half of the print, the coma aberration is higher in oculus than in cornea. It is therefore easily observed that the coma aberration is mainly caused by internal components (crystalline lens) (3). The spherical aberration is also higher in oculus than in cornea, indicating that spherical aberration is increased in the crystalline lens (4). Referring to the Landolt's ring simulation (5), it can be seen that the images are vertically distorted, which show the visual effect of the coma aberration. The image simulation is consisted of ocular HOA only, so it should be noted that this simulated image might not coincide with actual patient vision because the opacity of crystalline lens and subsequent light scattering affects the retinal image differently with ocular HOA.



Slit Lamp Anterior Image




Clinical Data ; Tokyo Medical and Dental University Hospital Faculty of Medicine
Editorial supervisor ; Hiroko Bissen-Miyajima, MD

Pseudophakic Eye with Toric IOL

This measurement shows in the upper half the IOL selection map, where it can be shown that the toric IOL is properly implanted and the corneal astigmatism is compensated by the implant. Reference values parameters are: Corneal Astigmatism (showing -1.79D@86), Ocular Astigmatism (showing -0.07D@91) and Internal Astigmatism (showing -1.72D@176). **1** In the lower half the PSF/MTF maps show that objective visual quality should be good as can be seen from the simulated PSF, that gives an indication of optical image quality, and from the MTF that gives an indication of contrast sensitivity **2**.

Customizable Print Layout



Example

2 analysis maps of your choice can be assembled to print layout on 1 page for printing and / or sending images to external systems, like IMAGENet.™

Specifications

Refractive Power Measuring Range

Sphere Range	-25D~+22D (0.01D/0.12D/0.25D steps) ※
Cylinder Range	0D~±10D (0.01D/0.12D/0.25D steps) ※
Axis Range	0°~180° (1°/5°steps)
Measurable Area	φ 8 mm (max.)
Measurable Minimum Pupil Diameter	φ 2mm

Corneal Curvature Measuring Range

Corneal Curvature Radius	5.00mm~10.00mm (0.01mm steps)
Corneal Refraction	67.5D~33.75D (0.01D/0.12D/0.25D steps) (Proviso: corneal refractive index = 1.3375)
Corneal Astigmatism	0D~±10D (0.01D/0.12D/0.25D steps)
Corneal Astigmatism Axial Angle	0°~180° (1°/5°steps)
Measurable Corneal Area	φ 0.8mm~ φ 9.2mm (Proviso Radius corneal curvature = 8mm)
Measurable PD Range	20~85mm (1mm step)
Export Output Terminal	USB (IN/OUT) 、RS232C (OUT) 、LAN (IN/OUT)

※Proviso:Sphere +Cylimder ≤ +22D, or Sphere+Cylimder ≥ -25D



Topcon Healthcare, Inc.
111 Bauer Drive, Oakland, NJ 07436, U.S.A.
Phone: +1-201-599-5100
www.topconhealthcare.com

IMPORTANT In order to obtain the best results with this instrument, please be sure to review all user instructions prior to operation.
Not all products, services, or offers are available in all markets. Contact your local distributor for country-specific information and availability.

CLASS 1 LASER PRODUCT
(IEC60825-1:2014)
PRODUIT LASER DE CLASSE 1
(CEI60825-1:2014)



TOPCON CORPORATION
75-1 Hasunuma-cho, Itabashi-ku, Tokyo 174-8580, JAPAN.
Phone: +81-(0)3-3558-2522/2502
Fax: +81-(0)3-3965-6898
www.topconhealthcare.jp

